



***Anthocharis cardamines* (Aurora)**



Vanessa atalanta (Almirante rojo)



Aglais io (Pavo real)



Papilio machaon (Macaón)

Castanea sativa (Castaño)

Prunus avium (Cerezo)

***Taraxacum officinale* (Diente de león)**

Genista florida (Piorno)

Betula sp. (Abedul)

Papaver rhoeas (Amapola)


Erica arborea (Brezo blanco)

***Erica australis* (Brezo rojo)**

Quercus robur (Roble común)



Phoenicurus ochruros (Colirrojo tizón)

A Common Cuckoo (Cuculus canorus) is perched on a thin, light-brown branch. The bird has a greyish-blue upper body with a yellowish-brown eye and a dark beak. Its underparts are white with dark horizontal stripes. The tail is dark with white spots. The background is a soft, out-of-focus green.

***Cuculus canorus* (Cuco)**

A Common House Martin (Hirundo rustica) is captured in flight over a body of water. The bird is shown from a side profile, with its wings fully extended, revealing the intricate structure of the feathers. Its body is dark blue-black, with a lighter, reddish-brown throat and breast. The background consists of a calm body of water reflecting the sky, with a rocky shoreline visible in the upper left. The overall scene is brightly lit, suggesting a sunny day.

Hirundo rustica (Golondrina)



***Coturnix coturnix* (Codorniz)**



Upupa epops (Abubilla)



Delichon urbicum (Avión común)



***Ciconia ciconia* (Cigüeña blanca)**



Milvus migrans (Milano negro)

Luscinia megarhynchos (Ruisseñor común)



Apus apus (Vencejo común)